



REVIEW ARTICLE

Anemia and iron deficiency in heart failure[☆]



Victor M. Gil^{a,*}, Jorge S. Ferreira^b

^a Serviço de Cardiologia, Hospital Fernando Fonseca, Amadora, Portugal

^b Serviço de Cardiologia, Hospital Santa Cruz – CHLO, Carnaxide, Portugal

Received 8 June 2013; accepted 25 June 2013

Available online 3 February 2014

KEYWORDS

Anemia;
Iron deficiency;
Heart failure;
Iron therapy

Abstract Heart failure is a common problem and a major cause of mortality, morbidity and impaired quality of life. Anemia is a frequent comorbidity in heart failure and further worsens prognosis and disability. Regardless of anemia status, iron deficiency is a common and usually unidentified problem in patients with heart failure. This article reviews the mechanisms, impact on outcomes and treatment of anemia and iron deficiency in patients with heart failure.

© 2013 Sociedade Portuguesa de Cardiologia Published by Elsevier España, S.L. All rights reserved.

PALAVRAS-CHAVE

Anemia;
Deficiência de ferro;
Insuficiência
cardíaca;
Terapêutica com
ferro

Anemia e deficiência de ferro na insuficiência cardíaca

Resumo A insuficiência cardíaca é uma patologia comum e uma causa importante de mortalidade, morbidade e deterioração da qualidade de vida. A anemia é uma comorbidade frequente na insuficiência cardíaca e agrava o seu prognóstico e capacidade funcional. Independentemente da presença ou não de anemia, a deficiência de ferro é um problema associado à insuficiência cardíaca muitas vezes não identificado. Este artigo revê os mecanismos, impacto prognóstico e tratamento da anemia e deficiência de ferro.

© 2013 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L. Todos os direitos reservados.

Case study

A 72-year-old woman with a history of poorly controlled hypertension under diuretic therapy for the previous 10 years was assessed for fatigue and dyspnea on minimal

exertion (NYHA class III). Her blood pressure was 150/85 mmHg and her heart rate was 88 bpm; she had faint crackles in the lung bases and ankle edema. The ECG showed voltage criteria for left ventricular hypertrophy. Her hemoglobin (Hb) level was 11.0 g/dl and her mean corpuscular volume was 85 fl, blood glucose 102 mg/dl, total cholesterol 190 mg/dl and creatinine 1 mg/dl. She was medicated with furosemide, a renin-angiotensin system inhibitor and a beta-blocker, and was referred for echocardiographic assessment.

What would did the patient's Hb of 11 g/dl play in her clinical setting?

[☆] Please cite this article as: Gil VM, Ferreira JS. Anemia e deficiência de ferro na insuficiência cardíaca. Rev Port Cardiol. 2014;33:39–44.

* Corresponding author.

E-mail address: victorgilmd@gmail.com (V.M. Gil).

Table 1 Factors associated with anemia in patients with heart failure.

1. Nutritional deficiency
2. Intestinal malabsorption
3. Hypovolemia with hemodilution
4. Renal failure
5. Hypothyroidism
6. Bleeding
7. Inflammation: increased serum cytokines (IL-6, TNF- α) and acute phase proteins (CRP)
8. Treatment with ACE inhibitors

ACE: angiotensin-converting enzyme; CRP: C-reactive protein; IL-6: interleukin-6; TNF- α : tumor necrosis factor alpha.

Introduction

The prevalence of heart failure (HF) is 1–2%, but may exceed 10% in individuals aged over 70.^{1,2} In Portugal, the prevalences of HF with systolic dysfunction and HF with preserved systolic function have been estimated at 1.3% and 1.7%, respectively, in a primary health care context.³

HF is associated with a marked decline in quality of life and high morbidity and mortality.⁴ Although treatment with renin-angiotensin-aldosterone system inhibitors and beta-blockers can be effective, one-year mortality is up to 20% and rehospitalizations for HF can reach 30%.⁵ However, the clinical course of HF is variable, due to the interaction of diverse demographic and clinical characteristics that affect prognosis, among them anemia and iron deficiency.^{6–8}

Anemia in heart failure

Anemia is a frequent comorbidity in HF. Its prevalence ranges between 4% and 55% depending on the study population and the cutoff values of Hb defined as the lower limit of normal.⁹ In a large cohort study published in 2003 of 12 065 patients with new-onset HF in 138 centers in the Canadian state of Alberta, the prevalence was 17%.¹⁰ It is more common in women, the elderly and individuals with renal failure.

Anemia is associated with increased morbidity and mortality, as well as significant reductions in functional capacity, which is already impaired by HF.^{6–8,11,12} In the largest observational registry published, anemia was an independent indicator of prognosis after correction for numerous confounding factors.¹²

There are various causes of anemia in association with HF (Table 1), including renal failure and hypothyroidism, and occasionally low levels of vitamin B₁₂ and folic acid.¹³ Another factor may be hemodilution, which can cause anemia due to expansion of plasma volume without any effective reduction in red blood cells. Angiotensin-converting enzyme inhibitors, which are commonly used in HF treatment, can be associated with low Hb, probably by suppression of erythropoietin, while inflammatory cytokines such as interleukin-1 and -6 and tumor necrosis factor alpha, which are elevated in severe HF, can reduce production of erythropoietin or increase resistance to its action. Finally, antiplatelet or anticoagulant therapy can cause gastrointestinal bleeding.¹⁴

Although all of the above factors can play a part, iron deficiency anemia (IDA) is the most common form in HF.

IDA associated with HF may be absolute or functional. The following mechanisms are presumed to be involved in the development of absolute IDA, in which iron stores are depleted as indicated by serum ferritin <30 $\mu\text{g}/\text{l}$ ¹⁵: (i) insufficient dietary iron supply, (ii) intestinal malabsorption, impaired duodenal iron transport, drug interactions (e.g. omeprazole), or food reducing absorption, and (iii) gastrointestinal bleeding.

In a 2006 study of the causes of anemia in HF patients, de Silva et al. found low iron or ferritin levels in 43% of patients but microcytosis in only 6%.¹⁶ However, Nanas et al. detected iron deficiency in bone marrow in their study population, despite normal serum iron, ferritin and erythropoietin.¹⁷ This may be explained by diversion of iron from the marrow to storage sites of the reticuloendothelial system, where it is unavailable for erythropoiesis despite normal or high iron and ferritin levels, as seen in chronic disease.¹⁸

An important factor is transferrin saturation (TSAT). Normal TSAT ranges between 20% and 50%; values below 20% are seen in IDA and inflammation anemia. In a study of 157 patients with chronic HF, TSAT <20% was associated with higher NYHA functional class, lower peak oxygen consumption and greater mortality in two-year follow-up.¹⁹

Functional IDA is defined as insufficient iron supply to meet demands, despite abundant iron stores, because iron is locked within cells of the reticuloendothelial system and is unavailable for cell metabolism. One mechanism in functional IDA appears to be related to hepcidin, a protein that is central to regulation of iron metabolism. Hepcidin levels are elevated in the initial stages of HF but fall as the disease progresses; low levels are an independent marker of worse prognosis.¹⁵ Hepcidin binds to ferroportin, the only protein able to export intracellular iron, which is thereby degraded, preventing iron from re-entering cells.¹⁵ This reduces iron absorption in the duodenum and causes iron to be retained in the reticuloendothelial system, reducing its concentration and availability in target tissues. Hepcidin also plays a part in erythropoiesis and in the innate immune response to pathogens.

Iron is essential for normal hematopoiesis; most of it is taken up by erythroblasts and reticulocytes for Hb synthesis. Iron deficiency results in resistance to hematopoietic growth factors (e.g. erythropoietin), and impairs the differentiation and maturation of all types of hematopoietic cells.¹⁵

Treatment of anemia in heart failure

There is no universally agreed definition of anemia. The World Health Organization considers anemia to exist when Hb levels are lower than 13 g/dl in men and 12 g/dl in women.²⁰ Others define it as hematocrit less than 35–39%.²¹ The target Hb level in patients with anemia is also not agreed, but 12 g/dl is regarded as safe.²²

The association of anemia with worse prognosis, functional capacity and quality of life in HF prompted efforts to correct this comorbidity. Besides blood transfusion, which is recommended for severe anemia,²³ there are two main types of treatment of anemia in HF: erythropoietic stimulating agents (ESAs) and iron supplementation.

Erythropoietic stimulating agents

Erythropoietin, which is produced in the kidneys in response to hypoxia, acts on the bone marrow to promote the survival and proliferation of erythroid precursor cells, increasing the production of red blood cells. Recombinant human erythropoietin (rHuEPO) and a more recent erythropoietin analog with a longer half-life, darbepoetin alfa, have been used in various trials to treat anemia associated with HF. The rationale for using ESAs is that although anemia in HF is multifactorial, it may represent an absolute or a relative deficiency or resistance to endogenous erythropoietin.

Silverberg et al. were the first to use ESAs in patients with anemia and HF, in an open randomized trial of 32 patients with moderate to severe HF and Hb 10–11.5 g/dL.²⁴ Treatment with rHuEPO and intravenous (IV) iron was followed by improvement in NYHA class, increased left ventricular ejection fraction and reductions in hospitalizations and need for diuretics. Several small-scale studies with rHuEPO or darbepoetin alfa have shown improvements in oxygen uptake and functional capacity, reduced BNP levels and improved quality of life.^{25–27} However, the results have been inconsistent. In a study by van Veldhuisen et al. of 165 patients with HF and Hb between 9.1 and 12.5 g/dL,²⁸ administration of darbepoetin alfa for a target value of 14 g/dL raised Hb levels and improved some quality of life indices as assessed by the Kansas City Cardiomyopathy Questionnaire, but no improvements were seen in the 6-minute walk test, NYHA functional class, or Minnesota Living with Heart Failure score.

A meta-analysis published in 2011 of 11 randomized clinical trials, nine placebo-controlled and five double-blinded, of a total of 794 patients with HF and anemia treated with ESAs showed improvements in symptoms and functional capacity and a reduction in clinical events.²⁹ It should be noted that in many of the small published iron supplementation was used in association with ESAs.

However, the latest evidence does not support the use of ESAs to treat anemia in HF. The lack of demonstrated benefit, and indeed the associated risks, were demonstrated in the Reduction of Events by Darbepoetin Alfa in Heart Failure (RED-HF) trial, which randomized 2278 patients with systolic HF and anemia to receive darbepoetin alfa or placebo, with a target Hb level of 13.0 g/dL.³⁰ Oral or IV iron was administered as tolerated if TSAT fell to less than 20% at any time. No significant differences were seen in the composite primary outcome of death from any cause or hospitalization for worsening heart failure in a mean follow-up of 28 months (50.7% in the darbepoetin alfa group and 49.5% in the placebo group (hazard ratio 1.01; 95% confidence interval [CI] 0.90–1.13) or in the incidence of stroke, although the latter was slightly higher in the darbepoetin alfa group (3.7% vs. 2.7%). However, thromboembolic events were more frequent in the darbepoetin alfa group (13.5% vs. 10%).

These results are similar to those of the Trial to Reduce Cardiovascular Events with Aranesp Therapy (TREAT), which compared darbepoetin alfa with placebo in 4028 patients with diabetes, chronic kidney disease, and anemia, a third of whom had HF.³¹ Higher rates of stroke, a thromboembolic events were seen in the darbepoetin alfa group.

Possible reasons for the adverse events seen with the use of ESAs include worsening of hypertension, elevated thromboembolic risk and increased release of endothelin.

Intravenous iron supplementation

The disappointing results reported for ESAs contrast with those described for IV iron supplementation alone in patients with HF and iron deficiency, even without manifest anemia.

The use of IV iron in the form of ferric carboxymaltose for the treatment of anemia of various causes was reviewed in a 2011 meta-analysis of 11 randomized trials and three cohort studies.³² A total of 2348 patients were treated with IV ferric carboxymaltose in randomized trials and 348 in cohort studies, 832 receiving oral iron, 762 receiving placebo and 384 receiving IV iron supplementation. Compared to oral iron (the most common comparator), IV ferric carboxymaltose increased mean Hb by 4.8 g/dL, ferritin by 163 µg/L and TSAT by 5.3%, and was significantly better than comparator in achieving the target Hb level (number needed to treat: 5.9). The incidence of serious adverse events was similar with IV ferric carboxymaltose and comparator, but gastrointestinal adverse events were significantly more frequent in patients treated with oral iron.

Six studies have been published on iron supplementation in HF, including two open non-controlled trials,^{33,34} one open controlled trial,³⁵ and three randomized placebo-controlled, double-blind trials^{36–38} (Table 2).

Improvements in symptoms, exercise capacity and quality of life were seen in parallel with rises in Hb levels in all these studies. In the Ferric Iron Sucrose in Heart Failure (FERRIC-H) trial, symptoms and exercise capacity also improved, although to a lesser extent, in non-anemic patients.³⁶ Improved oxygen consumption did not relate to absolute increases in Hb but to increased TSAT.

The multicenter randomized double-blind placebo-controlled Ferinject Assessment in Patients with Iron Deficiency and Chronic Heart Failure (FAIR-HF) trial was the largest.³⁸ Treatment with ferric carboxymaltose significantly improved the two main outcomes: 50% of patients randomized to ferric carboxymaltose reported at least moderate improvement on the Patient Global Assessment questionnaire, compared to 28% of the placebo group (odds ratio [OR] for improvement 2.51, 95% CI 1.75–3.61), and 47% of the treatment group achieved NYHA class I or II, compared to 30% in the placebo group (OR 2.40, 95% CI 1.55–3.71). Improvements were also seen in the 6-minute walk test and in quality of life scores, but prognosis was not affected (despite a tendency for fewer hospitalizations in the treatment group); the trial was not designed to assess the effect of iron supplementation on prognosis. The benefits of iron supplementation were seen in patients without as well as with anemia; this is slightly puzzling, but may be explained by the fact that only low-intensity exercise was tested, which correlates with iron levels in tissues. If maximum aerobic capacity had been tested, the non-anemic group would probably not have shown improvement, since this parameter depends on Hb concentrations rather than muscle oxidative capacity.³⁹

The question arises as to whether similar results could be obtained with oral iron. Absorption of oral iron(II) is blocked by hepcidin; moreover, gastrointestinal tolerance of oral iron drugs is generally poor, and they interact with various other medications, including proton pump inhibitors.⁴⁰ Furthermore, even if the iron is absorbed, its transport to the bone marrow is blocked by hepcidin, as in cases of anemia

Table 2 Studies on intravenous iron supplementation in patients with heart failure and anemia or iron deficiency.

Study	Design	n	Clinical context	Outcomes
Bolger et al. ³³	Open, non-controlled; IV iron sucrose	16	Hb \leq 12 g/dl; NYHA II/III	\uparrow Hb (1.4 g/dl); \uparrow QoL (MLHF); \uparrow 6-min WT (44 m)
Usmanov et al. ³⁴	Open, non-controlled; IV iron	32	Hb <11 g/dl; NYHA III/IV	\uparrow Hb (3–3.3 g/dl); \downarrow LV remodeling
Comin-Colet et al. ³⁵	Open; IV iron sucrose+EPO vs. no treatment	65	Hb \leq 12 g/dl; NYHA III/IV	\uparrow Hb (13.5 vs. 11.3 g/dl); \downarrow NT-proBNP; \downarrow hospitalization (25.9% vs. 76.3%)
FERRIC-HF ³⁶	Randomized, double-blind; IV iron sucrose vs. no treatment	35	FT <100 ng/ml or FT 100–300+TSAT <20%	\uparrow VO ₂ max 96 ml/min (p=0.08)
Tobli et al. ³⁷	Randomized, double-blind; IV iron sucrose vs. placebo	40	Hb <12.5 g/dl; FT <100 ng/ml; TSAT <20%; LVEF \leq 35%	\downarrow NT-proBNP; \uparrow QoL (MLHF); \uparrow 6-min WT
FAIR-HF ³⁸	Randomized, double-blind; ferric carboxymaltose vs. placebo	459	NYHA II/III; LVEF \leq 40%; FT<100 ng/ml or FT 100–299+TSAT <20%; Hb 9.5–13.5 g/dl	\uparrow PGA (50% vs. 28%); NYHA I/II (47% vs. 30%)

EPO: erythropoietin; FT: ferritin; Hb: hemoglobin; IV: intravenous; LV: left ventricular; LVEF: left ventricular ejection fraction; MLHF: Minnesota Living with Heart Failure score; PGA: Patient Global Assessment score; QoL: quality of life; TSAT: transferrin saturation; VO₂ max: peak oxygen uptake; 6-min WT: 6-minute walk test.

of chronic disease, which do not respond to oral iron. In IV formulations, iron in the form of iron(III) is bound to carbohydrate complexes, which are taken up from the bloodstream into the reticuloendothelial system within hours of administration. Most is incorporated into iron stores, but some binds to transferrin and reaches the bone marrow, thus overcoming blockade by hepcidin and resulting in underuse of this iron in tissue storage sites, and reduction of ferruliferous and increased risk of infection due to reduced cellular immunity and bacterial growth enhancement.⁴¹ However, the FAIR-HF trial did not show higher rates of infection or cardiovascular events. Unlike the iron dextran previously used, current IV iron preparations such as iron sucrose, ferric gluconate and ferric carboxymaltose are well tolerated, rarely causing hypersensitive reactions, and thus do not require a test dose.⁴²

A recent study shows improvements in previously low health-related quality of life after four weeks of treatment with ferric carboxymaltose irrespective of previous anemia status.⁴³

A health economic analysis of ferric carboxymaltose for HF patients with or without anemia in the UK showed the treatment to be cost-effective, below the threshold of 22 200–33 300 euros per quality-adjusted life year gained typically used by the UK National Institute for Health and Clinical Excellence, and robust in sensitivity.⁴⁴

Three clinical trials are currently underway comparing ferric carboxymaltose with placebo in patients with HF and iron deficiency: CONFIRM-HF (NCT01453608), with 300 patients, and iCHF (NCT01837082), with 100 patients, both ending in December 2013; and EFFECT-HF (NCT01394562), with 160 patients, due to end in January 2014. These trials are intended to determine the efficacy of ferric carboxymaltose in terms of improving functional capacity, and its safety. A combined analysis of the results of FAIR-HF and these trials will also show whether ferric carboxymaltose

reduces clinical events such as death and hospitalization for HF.

Guidelines for heart failure

There are no specific recommendations for the management of patients with HF and anemia or iron deficiency in the American or European guidelines (the latter adopted by the Portuguese Society of Cardiology).^{45,46} There is a general recommendation in the European guidelines for the management of anemia: "A standard diagnostic work-up should be undertaken in anemic patients. Correctable causes should be treated in the usual way."⁴⁵ It also states that IV ferric carboxymaltose may be considered for patients with iron deficiency.

Case study

Although not considered relevant at the initial assessment, the patient's Hb level of 11 g/dl will reduce her functional capacity and contribute to risk of morbidity and mortality. According to the European guidelines on HF, a diagnostic work-up of her anemia should have been performed, as well as medical therapy and echocardiographic assessment. Serum ferritin <100 ng/ml or 100–299 ng/ml associated with TSAT <20% indicates inflammation or chronic disease as the cause and should be corrected by iron supplementation, preferably parenterally.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Cleland JG, Khand A, Clark A. The heart failure epidemic: exactly how big is it? *Eur Heart J.* 2001;22:623–6.
- Mosterd A, Hoes AW. Clinical epidemiology of heart failure. *Heart.* 2007;93:1137–46.
- Ceia F, Fonseca C, Mota T, et al. Prevalence of chronic heart failure in Southwestern Europe: the EPICA study. *Eur J Heart Fail.* 2002;4:531–9.
- Stewart S, MacIntyre K, Hole DJ, et al. More 'malignant' than cancer? Five-year survival following a first admission for heart failure. *Eur J Heart Fail.* 2001;3:315–22.
- Jhund PS, MacIntyre K, Simpson CR, et al. Long-term trends in first hospitalization for heart failure and subsequent survival between 1986 and 2003: a population study of 5.1 million people. *Circulation.* 2009;119:515–23.
- Szachniewicz J, Petruk-Kowalczyk P, Majda J, et al. Anaemia is an independent predictor of poor outcome in patients with chronic heart failure. *Int J Cardiol.* 2003;90:303–8.
- Kosiborod M, Curtis JP, Wang Y, et al. Anemia and outcomes in patients with heart failure: a study from the National Heart Care Project. *Arch Intern Med.* 2005;165:2237–44.
- Komajda M, Anker SD, Charlesworth A, et al. The impact of new onset anemia on morbidity and mortality in chronic heart failure: results from COMET. *Eur Heart J.* 2006;27:1441–6.
- Felker GM, Adams Jr KF, Gattis WA, et al. Anemia as a risk factor and therapeutic target in heart failure. *J Am Coll Cardiol.* 2004;44:959–66.
- Ezekowitz JA, McAlister FA, Armstrong PW, et al. Anemia is common in heart failure and is associated with poor outcomes: insights from a cohort of 12 065 patients with new-onset heart failure. *Circulation.* 2003;107:223–5.
- Horwich TB, Fonarow GC, Hamilton MA, et al. Anemia is associated with worse symptoms, greater impairment in functional capacity and a significant increase in mortality in patients with advanced heart failure. *J Am Coll Cardiol.* 2002;39:1780–6.
- Kalra PR, Bolger AP, Francis DP, et al. Effect of anemia on exercise tolerance in chronic heart failure in men. *Am J Cardiol.* 2003;91:888–91.
- Anand S. Anemia and chronic heart failure: implications and treatment options. *J Am Coll Cardiol.* 2008;52:501–11.
- Murphy NF, McDonald K. Treatment of anaemia in chronic heart failure—optimal approach still unclear. *Eur Heart J.* 2007;28:2185–7.
- Jankowska EA, von Haehling S, Anker SD, et al. Iron deficiency and heart failure: diagnostic dilemmas and therapeutic perspectives. *Eur Heart J.* 2013;34:816–26.
- de Silva R, Rigby AS, Witte KK, et al. Anemia, renal dysfunction, and their interaction in patients with chronic heart failure. *Am J Cardiol.* 2006;98:391–8.
- Nanas JN, Matsouka C, Karageorgopoulos D, et al. Etiology of anemia in patients with advanced heart failure. *J Am Coll Cardiol.* 2006;48:2485–9.
- Weiss G, Goodnough LT. Anemia of chronic disease. *N Engl J Med.* 2005;352:1011–23.
- Okonko DO, Mandal AK, Missouri CG, et al. Disordered iron homeostasis in chronic heart failure: prevalence, predictors, and relation to anemia, exercise capacity, and survival. *J Am Coll Cardiol.* 2011;58:1241–51.
- Blanc B, Finch CA, Hallberg L, et al. Nutritional anaemias. Report of a WHO Scientific Group. WHO Tech Rep Ser. 1968;405:1–40.
- Tang WH, Yeo PS. Epidemiology of anemia in heart failure. *Heart Fail Clin.* 2010;6:271–8.
- Silverberg DS, Wexler D, Laina A, et al. The role of correction of anaemia in patients with congestive heart failure: a short review. *Eur J Heart Fail.* 2008;10:819–23.
- Tang YD, Klatz SD. Anemia in chronic heart failure: prevalence, etiology, clinical correlates, and treatment options. *Circulation.* 2006;113:2454–61.
- Silverberg DS, Wexler D, Sheps D, et al. The effect of correction of mild anemia in severe, resistant congestive heart failure using subcutaneous erythropoietin and intravenous iron. *J Am Coll Cardiol.* 2001;37:1775–80.
- Mancini DM, Katz SD, Lang CC, et al. Effect of erythropoietin on exercise capacity in patients with moderate to severe chronic heart failure. *Circulation.* 2003;107:294–9.
- Palazzuoli A, Silverberg D, Lovine F, et al. Erythropoietin improves anemia exercise tolerance and renal function and reduces B-type natriuretic peptide and hospitalization in patients with heart failure and anemia. *Am Heart J.* 2006;152, 1096.e9–e15.
- Ponikowski P, Anker SD, Szachniewicz J, et al. Effect of darbepoetin alfa on exercise tolerance in anemic patients with symptomatic chronic heart failure: a randomized, double-blind, placebo-controlled trial. *J Am Coll Cardiol.* 2007;20:753–62.
- van Veldhuisen DJ, Dickstein K, Cohen-Solal A, et al. Randomized, double-blind, placebo-controlled study to evaluate the effect of two dosing regimens of darbepoetin alfa in patients with heart failure and anaemia. *Eur Heart J.* 2007;28:2208–16.
- Kotecha D, Ngo K, Walters JA, et al. Erythropoietin as a treatment of anemia in heart failure: systematic review of randomized trials. *Am Heart J.* 2011;161:822–31.
- Swedberg K, Young JB, Anand IS, et al. Treatment of anemia with darbepoetin alfa in systolic heart failure. RED-HF Committees, RED-HF Investigators. *N Engl J Med.* 2013;368:1210–9.
- Pfeiffer T, Burdmann EA, Chen CY, et al. A trial of darbepoetin alfa in type 2 diabetes and chronic kidney disease. *N Engl J Med.* 2009;361:2019–32.
- Moore RA, Gaskell H, Rose P, et al. Meta-analysis of efficacy and safety of intravenous ferric carboxymaltose (Ferinject) from clinical trial reports and published trial data. *BMC Blood Disorders.* 2011;11:4.
- Bolger AP, Bartlett FR, Penston HS, et al. Intravenous iron alone for the treatment of anemia in patients with chronic heart failure. *J Am Coll Cardiol.* 2006;48:1225–7.
- Usmanov RI, Zueva EB, Silverberg DS, et al. Intravenous iron without erythropoietin for the treatment of iron deficiency anemia in patients with moderate to severe congestive heart failure and chronic kidney insufficiency. *J Nephrol.* 2008;21:236–42.
- Comin-Colet J, Ruiz S, Cladellas M, et al. A pilot evaluation of the long-term effect of combined therapy with intravenous iron sucrose and erythropoietin in elderly patients with advanced chronic heart failure and cardio-renal anemia syndrome: influence on neurohormonal activation and clinical outcomes. *J Card Fail.* 2009;15:727–35.
- Okonko DO, Grzeslo A, Witkowski T, et al. Effect of intravenous iron sucrose on exercise tolerance in anemic and non-anemic patients with symptomatic chronic heart failure and iron deficiency. FERRIC-HF: a randomized controlled, observer-blinded trial. *J Am Coll Cardiol.* 2008;51:103–12.
- Toblli JE, Lombraña A, Duarte P. Intravenous iron reduces NT-pro-brain natriuretic peptide in anemic patients with chronic

- heart failure and renal insufficiency. *J Am Coll Cardiol*. 2007;50:1657–65.
38. Anker SD, Colet JC, Filippatos G, et al. Trial Investigators. Ferric carboxymaltose in patients with heart failure and iron deficiency. *N Engl J Med*. 2009;361:2436–48.
 39. Davies KJ, Maguire JJ, Brooks GA, et al. Muscle mitochondrial bioenergetics, oxygen repletion, and work capacity during dietary iron deficiency and repletion. *Am J Physiol*. 1982;242:E418–27.
 40. Handelman GJ, Levin NW. Iron and anemia in human biology: a review of mechanisms. *Heart Fail Rev*. 2008;13:393–404.
 41. van Wyck D, Anderson J, Johnson K. Labile iron in parenteral iron formulations: a quantitative and comparative study. *Nephrol Dial Transplant*. 2004;19:561–5.
 42. Silverstein SB, Gilreath JA, Rodgers GM. Intravenous iron therapy: a summary of treatment options and review of guidelines. *J Pharm Pract*. 2008;21:431–43.
 43. Comin-Cole J, Lainscak M, Dickstein K, et al. The effect of intravenous ferric carboxymaltose on health-related quality of life in patients with chronic heart failure and iron deficiency: a subanalysis of the FAIR-HF study. *Eur Heart J*. 2013;34:30–8.
 44. Gutzwiller FS, Schwenkglens M, Blank PR, et al. Health economic assessment of ferric carboxymaltose in patients with iron deficiency and chronic heart failure based on the FAIR-HF trial: an analysis for the UK. *Eur J Heart Fail*. 2012;14:782–90.
 45. McMurray JJ, Adamopoulos S, Anker SD, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. *Eur Heart J*. 2012;33:1787–847.
 46. Yancy CW, Jessup M, Bozkurt B, et al. ACCF/AHA Guideline for the Management of Heart Failure. *J Am Coll Cardiol*. 2013, <http://dx.doi.org/10.1016/j.jacc.2013.05.019>.